

Barton R Branam, M.D.

University of Cincinnati Department of Orthopaedics and Sports Medicine Dr. Branam'sPost Operative Discharge Instructions Lower Extremity

*This is a standard copy of our discharge instructions. Discharge instructions may vary per patient depending on what procedures were performed. Please refer to your specific discharge instruction sheet.

Procedures:Knee arthroscopy, partial medial and/or lateral menisectomy, chondroplasty, excision of plica

Weight Bearing Status: as tolerated

Discharge Medications:

- _x_ Percocet 5/325 mg 1-2 tabs every 4-6 hours as needed for pain.
- _x_ Phenergan 25 mg 1 tablet every 6 hours as needed for nausea.
- _x_ Aspirin 325 mg daily starting tonight for 1 week post op.

May Remove Dressing On:3 days post op

- If you have steri-strips (white band-aid like strips directly over your incisions), please leave them on. These typically fall off in 1-2 weeks.

May Shower On:3 days post op

- Do not scrub your wound. Shower and let water runoff. Pat your wound dry.

- If after your shower, your wounds are dry then they can be left open to air. Keep your wound clean and dry. If fluid is coming out of your wound, then put a clean, dry dressing on your wound

- Please do not put any ointments or creams on your wound. This includes Neosporin, Bacitracin, etc.

- DO NOT TAKE A BATH OR SOAK YOUR WOUND. Call the office with any questions or concerns! Office Phone number: (513)475-8690

Please Make Follow-Up Appointment for 3-5 days post op.